

# HennesyMilner Logic and Bisimulation

Notes for the Course “Service Integration”

*Giuseppe De Giacomo*

Consider two transition system  $T = (A, S, s_0, \delta, F)$  and  $T' = (A, S', t_0, \delta, F')$  whose states we denote by  $s, s'$  and  $t, t'$  respectively.

Let  $L$  be the language formed by all the HennesyMilner Logic formulas. We define:

$$\sim_L = \{(s, t) \mid \forall \Phi \in L. T, s \models \Phi \text{ iff } T', t \models \Phi\}$$

and

$$\sim = \{(s, t) \mid \exists \text{ bisimulation } R \text{ s.t. } R(s, t)\}$$

Next we show that notably these two equivalence relations coincide!

**Theorem:**  $s \sim t$  implies  $s \sim_L t$ , i.e., If there exists a bisimulation between  $s, t$  then  $s, t$  satisfy (make true) the same formulas of HennesyMilner Logic.

**Proof:** By induction on the structure of the formulas. It suffices to consider only formula formed as follows:

$$\Phi \leftarrow Final \mid \Phi_1 \wedge \Phi_2 \mid \neg\Phi \mid \langle a \rangle \Phi$$

Indeed it is easy to see that  $\Phi_1 \vee \Phi_2 \equiv \neg(\neg\Phi_1 \wedge \neg\Phi_2)$  and  $[a]\Phi \equiv \neg\langle a \rangle\neg\Phi$ .

- **Atomic formulas** (*Final*) [*base case*]  $s \sim t$  implies  $s \in F$  iff  $t \in F'$  i.e.,  $T, s \models Final$  iff  $T', t \models Final$ .

- **Booleans** [*inductive cases*]

By induction hypothesis, we assume that for every  $s \sim t$  we have  $T, s \models \Phi_i$  iff  $T', t \models \Phi_i$ , for  $i = 1, 2$ . Then by  $T, s \models \Phi_1$  and  $T, s \models \Phi_2$  iff  $T', t \models \Phi_1$  and  $T', t \models \Phi_2$  hence, by definition we have  $T, s \models \Phi_1 \wedge \Phi_2$  iff  $T', t \models \Phi_1 \wedge \Phi_2$ .

Similarly for  $\neg\Phi$  (left as an easy exercise to the student).

- **Modal operators** [*another –the most interesting– inductive case*]

By induction hypothesis, we assume that for every  $ss \sim tt$  we have  $T, ss \models \Phi$  iff  $T', tt \models \Phi$ . Now consider that  $T, s \models \langle a \rangle \Phi$  iff there exists a transition  $s \rightarrow_a s'$  in  $T$  such that  $T, s' \models \Phi$ .

On the other hand since  $s \sim t$  there exists a transition  $t \rightarrow_a t'$  in  $T'$  such that  $s' \sim t'$ .

But then by induction hypothesis  $T, s' \models \Phi$  iff  $T', t' \models \Phi$ , and hence by definition  $T', t \models \langle a \rangle \Phi$ .  $\square$

**Theorem:**  $s \sim_L t$  implies  $s \sim t$ , i.e., If  $s, t$  satisfy (make true) the same formulas of HennessyMilner Logic, then there exists a bisimulation between  $s, t$ .

**Proof:** By coinduction. We show that  $\sim_L$  is a bisimulation, i.e., satisfy the following rules:

$$\begin{aligned} s \sim_L t \text{ implies} \\ s \in F \text{ iff } t \in F' \\ \text{if } s \rightarrow_a s' \text{ then } \exists t \rightarrow_a t' \text{ s.t. } s' \sim_L t' \\ \text{if } t \rightarrow_a t' \text{ then } \exists s \rightarrow_a s' \text{ s.t. } s' \sim_L t' \end{aligned}$$

• **Closure wrt the bisimulation rule**

– [local condition]

First, since  $s \sim_L t$  we have  $T, s \models Final$  iff  $T', s \models Final$ , but then we have  $s \in F$  iff  $t \in F'$ .

– [nonlocal condition]

We prove the rest by contradiction. Suppose that for some  $s, t$ , we have that  $s \sim_L t$ , and  $s \rightarrow_a s'$  but for all  $t \rightarrow_a t'$  we have  $s \not\sim_L t'$ . Then let  $\{t'_1, \dots, t'_n\} = \{t' \mid t \rightarrow_a t'\}^1$ . Notice since  $T, s \models \langle a \rangle True$  we have also  $T', t \models \langle a \rangle True$ , so  $n \geq 0$  above.

On the other hand, since  $s' \not\sim_L t_i$ , for each  $t'_i$  there is a formula  $\Phi_{t'_i}$  such that  $T', t'_i \models \Phi_{t'_i}$  but  $T, s \not\models \Phi_{t'_i}$ . Now consider the formula

$$[a](\bigvee_{i=1, \dots, n} \Phi_{t'_i})$$

Clearly  $T', t \models [a](\bigvee_{i=1, \dots, n} \Phi_{t'_i})$  but, since  $s \sim_L t$ , then also  $T, s \models [a](\bigvee_{i=1, \dots, n} \Phi_{t'_i})$ , which means that for all transitions  $s \rightarrow_a s''$  we must have  $T, s'' \models (\bigvee_{i=1, \dots, n} \Phi_{t'_i})$ , which is indeed false for  $s'' = s'$ . Contradiction.

Hence  $\sim_L$  itself is a bisimulation, so  $s \sim_L t$ , implies that  $s, t$  are bisimilar and hence  $s \sim t$ . □

---

<sup>1</sup>Here we assume that the transition systems are finite branching.